



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C12N 15/09, C12P 21/02, C07K 14/435	A2	(11) International Publication Number: WO 00/50586 (43) International Publication Date: 31 August 2000 (31.08.00)																																
<div style="display: flex; justify-content: space-between;"> <div style="width: 45%;"> (21) International Application Number: PCT/IB00/00219 (22) International Filing Date: 22 February 2000 (22.02.00) (30) Priority Data: 9904004.0 22 February 1999 (22.02.99) GB 9919369.0 16 August 1999 (16.08.99) GB </div> <div style="width: 50%;"> (81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). </div> </div>																																		
<div style="display: flex; justify-content: space-between;"> <div style="width: 45%;"> (71) Applicant (for all designated States except US): EUROPEAN MOLECULAR BIOLOGY LABORATORY [DE/DE]; Postfach 10 22 09, Meyerhofstrasse 1, D-69012 Heidelberg (DE). (72) Inventors; and (75) Inventors/Applicants (for US only): GEBAUER, Fatima [ES/DE]; Lerchenstrasse 28, D-69251 Gaiberg (DE). CORONA, Davide [IT/DE]; Brechtelstrasse 8-10, D-69126 Heidelberg (DE). BERGAMINI, Giovanna [IT/DE]; Römerstrasse 25, D-69115 Heidelberg (DE). BECKER, Peter [DE/DE]; Salzstrasse 9, D-82194 Gröbenzell (DE). HENTZE, Matthias [DE/DE]; Merianstrasse 5, D-69117 Heidelberg (DE). (74) Agent: HALLYBONE, Huw, George; Carpmals & Ransford, 43 Bloomsbury Square, London WC1A 2RA (GB). </div> <div style="width: 50%;"> Published <i>Without international search report and to be republished upon receipt of that report.</i> </div> </div>																																		
(54) Title: TRANSLATION SYSTEM																																		
(57) Abstract <p>The invention relates to a cell-free translation system using cell extracts, that reproduces the synergism that occurs during translation <i>in vivo</i> between the <i>cap</i> and the <i>poly(A)</i> tail structures of an mRNA molecule. Using this system, the amount of encoded protein produced from translation of an mRNA molecule is greater than the total of (a) the amount of the encoded protein that is produced under the same conditions when the ribonucleic acid template has a 5' cap but no 3' poly A tail, plus (b) the amount of the encoded protein that is produced under the same conditions when the ribonucleic acid template has a 3' poly A tail but no 5' cap.</p>																																		
<div style="text-align: center;"> Experiment E-872 Blow up (one set of data eliminated) c = cap, a = poly(A), c-a = cap and poly(A) Synergism = fold over the sum of c+a value </div> <div style="text-align: center;"> <table border="1" style="margin: 10px auto; border-collapse: collapse;"> <thead> <tr> <th>Extract</th> <th>Conditions</th> <th>c</th> <th>a</th> <th>c-a</th> <th>Synergism</th> </tr> </thead> <tbody> <tr> <td rowspan="2">Scott</td> <td rowspan="2">Our</td> <td>~200</td> <td>~200</td> <td>~1600</td> <td>8x</td> </tr> <tr> <td>~200</td> <td>~200</td> <td>~6800</td> <td>13x</td> </tr> <tr> <td rowspan="2">Scott</td> <td rowspan="2">Our</td> <td>~200</td> <td>~200</td> <td>~1000</td> <td>5x</td> </tr> <tr> <td>~200</td> <td>~200</td> <td>~4000</td> <td>11x</td> </tr> <tr> <td>Scott</td> <td>Our</td> <td>~200</td> <td>~200</td> <td>~200</td> <td>additive</td> </tr> </tbody> </table> </div>			Extract	Conditions	c	a	c-a	Synergism	Scott	Our	~200	~200	~1600	8x	~200	~200	~6800	13x	Scott	Our	~200	~200	~1000	5x	~200	~200	~4000	11x	Scott	Our	~200	~200	~200	additive
Extract	Conditions	c	a	c-a	Synergism																													
Scott	Our	~200	~200	~1600	8x																													
		~200	~200	~6800	13x																													
Scott	Our	~200	~200	~1000	5x																													
		~200	~200	~4000	11x																													
Scott	Our	~200	~200	~200	additive																													

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

TRANSLATION SYSTEM

The present invention relates to a cell-free translation system. In particular, the invention relates to a cell-free translation system derived from cell extracts, that reproduces the synergism that occurs during translation *in vivo* between the *cap* and the
5 *poly(A)* tail structures of the mRNA.

Translational control is an important mechanism for the regulation of gene expression during a variety of biological processes, including cellular metabolism (Hentze & Kuhn, (1996) P.N.A.S. USA 93: 8175-8182), cell differentiation (Ostareck *et al.*, 1997, Cell. 89: 597-606) and embryonic development (Wickens *et al.*, 1996, in Translation Control,
10 Eds. Hershey, J.W.B., Mathews, M.B. and Sonenberg, N. (Cold Spring Harbour Lab. Press, Plainview, NY, pp. 411-450). Failure to regulate translation properly often leads to disease.

In vivo, two modifications in the mRNA, namely a *cap* structure at the 5' end and a *poly(A)* tail at the 3' end, act synergistically to promote mRNA translation (Gallie, 1991,
15 Genes Dev. 5: 2108-2116). To date, research into the function of those structures and the synergism that occurs between them has been restricted by the lack of *in vitro* systems that accurately recapitulate these features.

Recently, a cell-free translation system that reproduces this synergism has been obtained from the unicellular eukaryote *Saccharomyces cerevisiae* (Iizuka *et al.*, 1994,
20 Mol. Cell. Biol. 17: 7322-7330). Capped mRNAs were found to be translated 50-fold more efficiently than uncapped mRNAs in yeast lysate. Polyadenylated RNAs were found to be translated 140-fold more efficiently than non-polyadenylated RNAs. mRNAs containing both a 5' *cap* and a 3' *poly(A)* tail were found to be translated several hundred-fold more efficiently than mRNAs lacking both of these terminal sequences.

25 A cell-free translation system derived from animal cells that is capable of reproducing this synergism would be of great value for a number of reasons. Firstly, this would facilitate and enhance the validity of basic research into the factors and mechanisms involved in translational control in higher eukaryotes, including the regulation of translation by cis- and trans- acting factors, the molecular basis of synergism, mRNA

stability and polyadenylation. Second, species-specific differences in translational mechanisms could be evaluated. Such a system would also allow the improvement of large-scale protein synthesis, both with respect to the efficiency of translation and to the fidelity of post-translational modification.

5 Summary of the Invention

According to a first aspect of the present invention, there is provided a method for the *in vitro* translation of a ribonucleic acid template, said ribonucleic acid having both a 5' cap and a 3' poly A tail, said method comprising incubating a cell extract of a multicellular eukaryote with said ribonucleic acid template under conditions such that translation of the RNA template to produce its encoded protein by one or more components in the cell extract occurs and the amount of the encoded protein thus produced is greater than the total of (a) the amount of the encoded protein that is produced under said conditions when the ribonucleic acid template has a 5' cap but no 3' poly A tail, plus (b) the amount of the encoded protein that is produced under said conditions when the ribonucleic acid template has a 3' poly A tail but no 5' cap.

It has been found, surprisingly, that by combining a dechoriation procedure that has been previously used for the study of chromatin assembly (Becker *et al.*, 1994) with homogenisation and centrifugation conditions used to study translation (Scott *et al.*, 1979) a protocol results that allows the preparation of cell extract with greatly improved properties over conventionally known extracts for use in *in vitro* translation procedures.

The cell extract may be an animal cell extract, such as a mammalian cell, or an insect cell. Specific examples include cell extracts of *Drosophila*, *C. elegans*, rodent, rabbit (for example, reticulocyte cells), human (for example HeLa cells) and primate cells. The cells may be embryonic or adult or from a particular tissue such as the ovaries. Preferably, the cell extract is a *Drosophila* cell extract, most preferably a *Drosophila* embryo or ovary cell extract.

The extract exhibits properties that were previously unforeseen, such as the capability to reproduce the synergism that occurs during translation *in vivo* between the cap and the poly(A) tail structures of an mRNA molecule, and the capability to perform mRNA polyadenylation. In addition, the procedure is simple and allows the preparation of large

amounts of extract at low cost. This makes the system useful in both basic research and in commercial applications as is discussed in more detail below.

RNA templates suitable for use in the methods of *in vitro* translation of the invention
5 may be recombinant or can be purified native templates. In preferred aspects of the invention, the poly(A) tail of the RNA template can be encoded in a DNA coding sequence which can be transcribed to generate an RNA template with a poly(A) tail of defined length. An alternative method of generating the poly(A) tail is the use of a poly(A) polymerase in an *in vitro* reaction to add the tail to the template as a post-
10 transcriptional modification.

The 5' cap portion may be added co-transcriptionally to the RNA template by the RNA polymerase. A suitable protocol can be found in "Protocol and Applications Guide"; Promega, 2nd edition, p62. If the template is a purified native RNA template, the cap
15 structure will already be in place.

According to a second aspect of the invention there is provided a method for the preparation of a *Drosophila* embryo extract comprising the steps of a) dechorionating *Drosophila* embryos in an aqueous isotonic buffer comprising detergent and bleach; b)
20 washing the embryos; c) homogenizing the embryos to produce a homogenate; d) centrifuging the homogenate; and e) recovering non-pelleted material from the centrifuged homogenate. This cell extract is particularly suited for use in the method of the first aspect of the invention.

The age of the embryo is not crucial to the method of the invention, although embryos
25 of between the ages of 1 hour and 12 hours of development are preferred.

Before preparation of extract, the embryos should preferably be collected in a sieve and washed extensively with water to remove debris from the culture medium in which they have been laid. For preparation of extract, the embryos are washed with an aqueous buffer, preferably with fresh EW buffer (0.7% NaCl, 0.04% Triton X-100), and poured
30 into a suitable vessel. An excess of buffer is then added and the embryos allowed to settle. At this stage, floating embryos should be discarded. For dechoriation, fresh buffer should be added and the embryo suspension agitated, preferably using a spinning

magnet or other suitable device. In all steps for the preparation of cell extract, ethanol-based buffers should be avoided.

For dechoriation, bleach is added to the embryo suspension, preferably at a concentration of about 3% by volume and the solution incubated for an appropriate
5 time. Preferably, incubation is for between 1 and 9 minutes, most preferably 3 minutes.

Dechoriation may be carried out at a temperature of between 18 and 37°C. Preferably, dechoriation is carried out at about 25°C.

After dechoriation, embryos are preferably transferred to a sieve and washed extensively with water. Dechorionated embryos are then transferred to a suitable vessel
10 (such as a beaker or cylinder) and buffer such as DE buffer (10mM HEPES pH 7.4, 5mM DTT) added. Floating embryos are discarded. Preferably, about 1 volume (with respect to the volume of settled embryos) of fresh buffer containing protease inhibitors is added to prevent proteolysis of components of the extract.

Embryos are then homogenised. Preferably, homogenisation is carried out at about 4°C.
15 Most preferably, a Potter-Elvehjem homogeniser is used at about 1500 rpm (about 20 strokes). The homogenate should preferably be kept on ice to minimise proteolysis.

The homogenate is preferably centrifuged, ideally in an ultracentrifuge at 24,000 rpm (40,000g)/TLS-55 rotor/4°C/20 minutes. The clear interphase is taken, ideally by puncturing the tube with a syringe, and is transferred to a Falcon tube.

20 Glycerol may then be added to the extract, preferably to a 10% final concentration, and the extract aliquoted, frozen in liquid nitrogen and kept at -70°C for future use.

Drosophila ovaries may also be used to produce a cell extract. The procedure is as follows: adult female flies are manually dissected to obtain the ovaries, which should be placed in an eppendorf tube on ice in an aqueous medium, for example PBS. Ovaries
25 are then allowed to settle, before transferring the ovaries slowly from an isotonic medium to a hypotonic medium. For example, the ovaries can be washed twice with around 12 volumes of a 1:1 mix of PBS:DEI buffer (DEI = 10mM HEPES pH 7.4, 5mM DTT, 1 x COMPLETE-Protease inhibitors from Boehringer Mannheim cat#1697498), and twice with 12 volumes of DEI. After washing, all the buffer is

removed and the ovaries manually homogenised using a plastic pestle. The homogenate can then be spun and the extract collected as described above for embryo extract.

The invention also provides a method for the preparation of a mammalian cell extract comprising the steps of a) collecting cells by centrifugation; b) washing the cells; c) 5 resuspending the cells; d) homogenising the cells to produce a homogenate; e) centrifuging the homogenate and f) recovering non-pelleted material from the centrifuged homogenate. This cell extract is well-suited to use in the method of the first aspect of the invention.

Before preparation of extract, the cells should be grown in culture, preferably at 10 exponential growth. Suspension cultures grown in, for example Jocklik's medium supplemented with 5% newborn bovine serum at 37°C, are suitable. The cells may be collected from culture by centrifugation, for example, by harvesting at 700g for 15 minutes. For preparation of extract, the cells should be washed with phosphate buffered saline at 4°C, before resuspension in a suitable ice cold hypotonic buffer such as a 15 buffer containing containing Hepes 10 mM. PH 7.6. KOAc 10 mM. Mg(OAc)₂ 0.5 mM and Dithiothreitol 5 mM. Reduced amounts of DTT are not recommended and addition of protease inhibitors is optional.

After a short period on ice, for between 2 and 10 minutes, the cells are homogenised. Preferably, homogenisation is carried out at about 4°C. Most preferably, the cells are 20 homogenised using between 10 and 30 strokes of a Dounce homogeniser (pestle type B). The homogenate should preferably be kept on ice to avoid proteolysis.

The homogenate is then centrifuged, ideally at 14000g for 5 minutes at 4°C. A longer centrifugation is not recommended. Supernatant fluid is divided into aliquots frozen in liquid nitrogen and stored at -80° C. Addition of glycerol is not necessary.

25 According to a third aspect of the invention there is provided the extract of *Drosophila* embryos or ovaries, or of mammalian cells, produced according to the methods of the second aspect of the invention.

According to a fourth aspect of the invention there is provided a method for the *in vitro* translation of a ribonucleic acid template, comprising the steps of adding a ribonucleic

acid template to a translation mix in the presence of the embryo, ovary or mammalian cell extract of the third aspect of the invention to form a reaction mix and incubating the reaction mix for at least 90 minutes at between 18°C and 28°C. Preferably, the reaction mix is incubated for at least 90 minutes at 25°C.

- 5 By reaction mix is meant a solution of components necessary for *in vitro* translation to occur. Such components may include any of the following: spermidine, amino acids, creatine phosphate, creatine kinase, dithiothreitol (DTT), buffer, $Mg(OAc)_2$, KOAc, tRNA, cell extract and RNA template. This reaction mix should preferably be prepared fresh. All the components of the reaction mix, except the creatine kinase, should be
0 dissolved in distilled water to their required concentrations.

Preferably, the buffer is HEPES buffer, although any buffer with a low concentration of salt (below 10mM) may be used. HEPES buffer is most preferably used at a final concentration of between 16 and 24mM. As used herein, the term "final concentration" means the concentration of the component that is present in the *in vitro* translation mix.

- 5 These concentrations are optimal for the translation reaction to take place.

Preferably, said tRNA is calf liver tRNA, at a final concentration of below 150µg/ml, most preferably about 100µg/ml.

- For *Drosophila* ovary or embryo extract, the final concentration of spermidine is preferably about 0.1mM. For amino acids, final concentration is about 60µM. For
10 creatine phosphate, the concentration can be in the range 17-23mM. It is shown herein (see Figure 6) that it is preferable to use fresh or newly thawed creatine phosphate. The level of translation doubles when fresh creatine phosphate is used. Creatine kinase should be used at a concentration of about 0.08mg/ml.

- The concentrations of DTT, $Mg(OAc)_2$ and KOAc should be optimised for each
15 individual mRNA template, as will be appreciated by those skilled in the art. For example, for translation of M1414WT (bgal) (see below), ideal concentrations are DTT, 1.2mM; $Mg(OAc)_2$, 0.6mM; and KOAc, 60mM. For c-Luc-a, ideal concentrations are DTT, 0mM; $Mg(OAc)_2$, 0.4mM; and KOAc, 30mM.

However, final concentration of $\text{Mg}(\text{OAc})_2$ can be in the range 0-3mM. KOAc can range between 0 and 200mM. DTT can range between 0-4mM.

Embryo or ovary extract should preferably be used at a concentration of about 40% by volume.

- 5 For mammalian cell extract, the final concentration of spermidine is preferably about 0.1mM. For amino acids, final concentration is about 100 μ M. For creatine phosphate, the concentration can be in the range 17-23mM. Creatine kinase should be used at a concentration of about 0.05mg/ml. As stated above for *Drosophila* extract, the concentrations of $\text{Mg}(\text{OAc})_2$ and KOAc should be optimised for each individual mRNA
- 10 template, as will be appreciated by those skilled in the art. For example, suitable concentrations may be along the following lines; $\text{Mg}(\text{OAc})_2$, 2.5mM; KOAc, 50mM.

However, final concentration of $\text{Mg}(\text{OAc})_2$ can be in the range 0-3mM. KOAc can range between 0 and 200mM. DTT can range between 0-4mM.

- Mammalian cell extract should preferably be used at a concentration of about 40% by
- 15 volume.

- Optimal RNA template concentration will vary depending upon the RNA type. The skilled artisan will appreciate that optimisation of the system may be easily performed for a particular RNA species of choice using routine procedures. For example, for the species c-luc-a (coding for firefly luciferase) a concentration of about 3.2ng/ μ l is
- 20 optimal using *Drosophila* embryo extract. The mRNA template should preferably be capped and should contain a *poly(A)* tail of more than 31 adenine nucleotides (more preferably 70-100 adenines).

- For each mRNA species, preliminary reactions should be performed so as to obtain a curve of translation efficiency with differing amounts of mRNA. An amount of mRNA
- 25 should be used in the linear range of translation. In the examples described herein, test reactions were performed with M1414WT (bgaI) [this codes for the β -galactosidase enzyme and contains 5' and 3' UTR sequences from *Drosophila oskar* mRNA], c-Luc-a and Luc mRNA (Promega).

Reaction mix prepared in this fashion contributes to the advantageous features of the method of this aspect of the invention, namely reproducing the synergy between the 5' *cap* and 3' *poly(A)* tail of the RNA molecule for translation.

- Additionally to this synergy, the level of translation is enhanced, so leading to a greater
- 5 production of protein. The fidelity of post-translational modification appears also to be retained, differentiating the system of the present invention over that described previously in yeast (Iizuka *et al.*, 1994). Although yeasts do possess enzymes that are capable of effecting the post-translational modification of proteins, the type and extent of modification tends to differ from that found in higher eukaryotes. Consequently, the
- 10 *Drosophila* system of the present invention generates proteins modified similarly to the state in which they are found *in vivo*.

The incubation step is preferably performed for sufficient time to allow the translation reaction to proceed to the extent desired. Preferably, the incubation step is performed for at least about 90 minutes.

- 15 Ranges of suitable concentrations of components of the reaction mix for *Drosophila* extract, along with optimal concentrations, are given below in Table 1. These concentrations were optimised for luc and c-luc-a mRNA species, coding for the firefly luciferase enzyme.

TABLE 1

	Range tested	Optimal	RNA	Observations
Creatine phosphate	0-168mM	17-23mM	Luc	Use fresh (thaw only once)
Creatine kinase	0-1.68 mg/ml	0.08 mg/ml	Luc	
KOAc	0-200mM	74mM	Luc	Should be optimised for each RNA template
Mg(OAc) ₂	0-3mM	0.3mM	Luc	Should be optimised for each RNA template

Spermidine	0-1.8mM	0.1mM	c-Luc-a	
Amino acids	0-100 μ M	60 μ M	c-Luc-a	
DTT	0-4mM	0mM	c-Luc-a	Should be optimised for each RNA template
tRNA	0-150 μ g/ml	100 μ g/ml	Luc	
Temperature	18-37°C	25°C	c-Luc-a	

Ideally, the *in vitro* translation reaction is performed in a final volume of about 12.5 microlitres. Below, in Table 2, the volume of stock concentrations is given in order to make up a reaction volume of 12.5 microlitres.

5 TABLE 2

- Final volume = 12.5 μ l
- Mix per reaction:

	Volume (μ l)	Final conc.
2.5mM spermidine	0.5	0.1mM
1mM amino acids (Promega)	0.75	60 μ M
1 M creatine phosphate (fresh or newly thawed)	0.21	16.8mM
10 μ g/ μ l creatine kinase	0.1	80 ng/ μ l
0.1 M DTT	0.15	1.2mM
1 M HEPES pH 7.4	0.3	24mM
4.13mM Mg(OAc) ₂	1.8	0.6mM

1 M KOAc	0.75	60mM
1.1 $\mu\text{g}/\mu\text{l}$ calf liver tRNA	1.1	100ng/ μl
Embryo extract	5	40%
RNA-template	1	3.2ng/ μl
H ₂ O	to 12.5 μl final	

For HeLa cell extract, the reaction mix may be as follows:

Untreated extract	40% of the final volume
Hepes buffer, pH 7.6	16 mM
KOAc	50 mM
Mg(Oac) ₂	2.5 mM
amino acid mixture	100 μM
spermidine	0.1 mM
ATP	0.8 mM
GTP	0.1 mM
Creatine phosphate	20 μM
Creatine kinase	50 ng/ μl
Calf liver tRNA	100 ng/ μl
RNA template	1 ng/ μl

These optimal values were obtained for this translation assay by testing a range of components concentrations with a capped and polyadenylated mRNA expressing the bacterial enzyme chloramphenicol acetyltransferase (CAT). The amount of the CAT

protein produced was measured using a colorimetric enzyme immunoassay (CAT ELISA, Böhrringer Mannheim).

According to a fifth aspect of the invention there is provided the use of cell extract according to the third aspect of the invention in a method of *in vitro* translation of a
5 ribonucleic acid template, such as the methods of the first and fourth aspects of the invention. The method reproduces the synergism exhibited *in vivo* between the 5' cap and 3' poly(A) tail of the ribonucleic acid template.

According to a sixth aspect of the invention there is provided the use of cell extract according to the third aspect of the invention or method of first or the fourth aspects of
10 the invention to screen for compounds that either a) decrease the translation, polyadenylation and/or stability of all mRNA species, or of specific mRNAs; b) potentiate the translation, polyadenylation and/or stability of all mRNA species, or of specific mRNAs; or c) decrease or potentiate the synergism between the 5' cap and the 3' poly(A) tail of the mRNA. Suitable compounds may interact (either directly or
15 indirectly) with only one of the 5' cap or poly(A) structures. Preferably, compounds identified in such a screen have an effect on translational control in higher eukaryotes, in particular insects. This screening method has advantages over prior art methods because the synergy exhibited between cap and poly(A) structures mimics the features of *in vivo* translation and allows the screening in an *in vitro* system for modifiers
20 effecting translation that are significantly more likely to function *in vivo*.

According to a seventh aspect of the invention there is provided a compound identified by a screen according to the sixth aspect of the invention. Suitable compounds may be natural or synthetic, for example, an aptamer, or a peptide and may be useful for therapy of diseases and for the control of plagues and diseases caused by insects, for example
25 malaria and plant pestilence. For example, the compound may that interact with 5' cap or 3' poly(A) structures of an mRNA template or affect in some way the synergism between the 5' cap and 3' poly (A) tail of an mRNA template. In some instances, compounds may be specific for a certain group of mRNA templates, or for mRNA templates within a certain species. It is envisaged that such compounds will prove
30 useful in the control of plagues and diseases caused by insects, for example malaria and plant pestilence, by interfering with the expression of certain essential insect proteins.

According to an eighth aspect of the invention there is provided the use of the cell extract according to the third aspect of the invention or method of the first or fourth aspects of the invention in research into the regulation and function of post-transcriptional mRNA modification or post-translational protein modification.

- 5 After the identification of the polypeptides that are responsible for the synergism between the 5' cap and the 3' poly(A) structures of the mRNA, it is possible to study the productive interactions between them and to define the domains and amino acid residues that are responsible for these interactions. It will, then, become possible to design chimeric molecules that comprise the effector domains from these polypeptides
- 10 fused to other polypeptides of interest. Upon interaction of the polypeptides of interest, the effector domains would promote the translation of a reporter mRNA, such as luciferase. The level or degree of interaction would be quantified by measuring the amount of light that is produced. In order to aid the study of interactions of interest, appropriate components such as *Drosophila* embryo extract or HeLa cell extract,
- 15 reporter mRNA and reaction mix may be provided in the form of a kit.

According to a ninth aspect of the invention there is provided a kit for the analysis of *in vitro* translation, polyadenylation and/or stability of ribonucleic acid. The kit should contain cell extract according to the third aspect of the invention along with stock solutions of all the other components of the translation mix in appropriate quantities.

- 20 The kit will therefore in addition comprise any one or all of spermidine, amino acids, creatine phosphate, creatine kinase, optionally dithiothreitol (DTT), buffer, $Mg(OAc)_2$, KOAc, tRNA, and RNA template.

- As noted above, preferably, the buffer is HEPES buffer, the tRNA is calf liver tRNA and ideally, the creatine phosphate is made fresh. However, creatine phosphate may be
- 25 supplied in powdered form in the kit for dissolution in distilled water as and when required for use.

- Generally, an RNA template will not be supplied as part of the kit, since in the main part, it is envisaged that the kit will be purchased for the purposes of experimental research. However, a control RNA template may be included in the kit so that a user can
- 30 ensure that the translation reaction is proceeding appropriately. For example,

comparison with a control RNA template may be used to optimise the concentration of test RNA, DTT, $\text{Mg}(\text{OAc})_2$ and KOAc that is to be used.

The kit preferably also contains appropriate instructions to enable a user to perform the *in vitro* translation reaction appropriately.

- 5 The invention will now be described in detail with particular reference to a Luc mRNA template. As will be clear to the person of skill in the art, variations from the described protocol may be made without departing from the scope of the invention.

All references cited herein are hereby incorporated by reference in their entireties.

BRIEF DESCRIPTION OF THE FIGURES

- 10 Figure 1 shows comparative results of translation levels and levels of synergy found between 5' *cap* and 3' *poly(A)* tail of an RNA template.

Figure 2 is an expanded scale of the results of Figure 1.

Figure 3 shows optimisation of Mg^{2+} concentration.

Figure 4 shows optimisation of K^+ concentration.

- 15 Figure 5 shows optimisation of creatine kinase concentration.

Figure 6 shows how percentage translation is improved by using fresh creatine phosphate.

Figure 7 shows optimisation of creatine phosphate concentration.

Figure 8 shows optimisation of spermidine concentration.

- 20 Figure 9 shows optimisation of tRNA concentration.

Figure 10 shows optimisation of temperature, amino acid concentration and DTT concentration using c-luc-a RNA.

Figure 11 shows the time course of translation of CAT mRNAs using the HeLa cell extract.

Figure 12 shows Northern blot analysis of the transcripts during the translation assay. Bars represent radioactive intensity of each messenger measured by a phosphoimager. Graph (A) shows the four messengers at different times, from 0 to 150 minutes. Graph (B) shows the Cap and the Cap-pA transcripts from 0 to 90 min.

- 5 Figure 13 shows time course of translation of the Luc messengers. Efficiency of translation is represented by light emission measured by a luminometer. Graph (B) shows the data represented in Graph (A) with the y axis maximum at 20000 luminescence instead of 550000.

EXAMPLES

10 Example 1: Preparation of extracts from *Drosophila* embryos

Extracts from 1h30', 3h, 6h and 12h embryos were prepared with similar results.

- Embryos were laid by 2-3 day old adult *Drosophila* flies on agar-apple juice plates (2.9% Agar; 30% Apple Juice; 4.4% Rubensirup; 0.25% Nipagin). The plates, spread with some fly food (220ml deionized water, 1.4ml Propionic acid; 150g dry yeast), were
15 left overnight for embryo laying). Embryos from 16 agar-apple juice plates were collected in a pile of sieves (the first with a cut-off size of an adult fly, the second with a cut-off size of a fly appendix [e.g.: a leg, a head or an antenna], the last with a cut-off size of a single embryo) and washed extensively (~5-10 minutes) with tap water to remove debris coming from the plates.

- 20 Embryos were subsequently washed with freshly prepared isotonic EW buffer (0.7% NaCl, 0.04% TritonX-100). To do this, embryos were transferred to a 500ml cylinder containing EW buffer, and were allowed to settle for ~3-5 minutes (roughly the time needed for 90% of the embryos to pellet by gravity) and were then washed twice with 500ml EW Buffer. Floating embryos were eliminated by suction.

- 25 Embryos were dechorionated in the 500 ml cylinder at room temperature (~20-25°C) with 260ml of EWB (0.7% NaCl, 0.04% TritonX-100, 3% Sodium Hypochlorite) for 3 minutes under vigorous agitation provided by a magnetic stirrer. Sodium Hypochlorite was from Sigma, Thomas Chemikalien, or U.S. CHLOROX. No significant change in the translation efficiency of the extract was noted for different bleach types.

Dechorionated embryos were transferred back to the sieves, and were vigorously and extensively washed with tap water (by flushing a strong stream of water for about 5-10 minutes). Washed embryos were settled by gravity twice with 100 ml of DE buffer (10mM HEPES pH 7.4, 5mM DTT) in a 100ml cylinder. Floating embryos were discarded, and an equivalent of one volume (with respect to the settled embryos) of DEI buffer (10mM HEPES pH 7.4, 5mM DTT, 1x COMPLETE-Protease Inhibitors from Boehringer Mannheim cat# 1697498) was added.

Embryos in DEI buffer were homogenised in a cold room (~4°C) by 20 strokes of a Potter-Elvehjem homogeniser at 1500 rpm and the homogenate was kept on ice. The homogenate was spun in a table-top ultracentrifuge (Beckmann) at 24000 rpm (40000x g) in a TLS-55 rotor at 4 C for 20 minutes. The clear aqueous interphase was taken by puncturing the tube with a syringe, and was transferred to a Falcon tube. Glycerol was added to 10% final, and the extract was aliquoted, flash-frozen in liquid nitrogen and stored at -70°C.

15 Example 2:

2.1 *In vitro* translation

A typical experiment is shown below.

The reaction is performed in a final volume of 12.5 µl, containing the following mix (a):

20		<u>volume (µl)</u>	<u>Final conc.</u>	
	2.5 mM spermidine	0.5	0.1 mM	
	1mM amino acids	0.75	60 µM	
	1 M creatine phosphate (fresh or newly thawed)	0.21	16.8 mM	
25	10 µg/µl creatine Kinase	0.1	80 ng/µl	
	0.1 M DTT	Y	0 - 1.2 mM	(b) (*)
	1 M HEPES pH 7.4	0.3	24 mM	
	4.13 mM Mg(OAc) ₂	Z	0.3 - 0.6 mM	(b) (*)
	1 M KOAc	W	30 - 80 mM	(b) (*)
30	1.1 µg/µl calf liver tRNA	1.14	100 ng/µl	

embryo extract	5	40%
RNA template	X	(c)
H ₂ O	to 12.5 µl final	

The reaction was incubated at 25°C for 90 min.

5 (a) A master mix should be prepared for as many samples as it is required.

(b) Should be optimised for each mRNA template.

(c) The mRNA template should be capped and should contain a poly(A) tail of more than 31 A's (usually 70-100 A's), although mRNAs without a poly(A) tail are also translated albeit at lower efficiency. A curve of translation with different amounts of
10 RNA should be performed to use an amount in the linear range of translation.

(*) Concentrations (mM) optimal for:

	<u>RNA</u>	<u>DTT</u>	<u>Mg(OAc)₂</u>	<u>KOAc</u>
	M1414WT (bgal)	1.2	0.6	60
	c-Luc-a	0	0.4	30
15	mLuc-a	0	0.6	80
	c-CAT-a	1.2	0.6	60

M1414WT RNA encodes for the β-galactosidase enzyme (bgal) and contains 5' and 3' UTR sequences from *Drosophila oskar* mRNA. *c-Luc-a* encodes for the firefly luciferase enzyme. *mLuc-a* encodes for the firefly luciferase enzyme, and contains the
10 5' UTR of *Drosophila msl-2* mRNA. *c-CAT-a* encodes for the chloramphenicol acetyl transferase enzyme. All RNAs are capped and polyadenylated.

Spermidine is from SIGMA, cat# S-0381.

Amino acids were obtained from Promega (cat #L4461), or can be self-made from a SIGMA kit (cat# LAA-21).

15 Creatine Kinase is from Boehringer Mannheim (cat# 126969). The stock solution is prepared at 10 mg/ml in 50% glycerol, 20 mM HEPES-KOH pH 7.4.

Using this protocol, together with the extract described in Example 1, high levels of translation and synergy can be obtained; compared to other methods (see Figure 1).

2.2 Optimisations

Ranges of suitable concentrations, along with optimal concentrations, are given below.

- 5 The method of Example 1 was followed (not that of Scott *et al*), and one concentration value was varied for each experiment. Luc RNA from Promega was used for these experiments, except where otherwise stated.

Figure 3 shows optimisation of Mg^{2+} concentration. Optimum concentration was found to be 0.3mM. Figure 4 shows optimisation of K^{+} concentration. Optimum concentration
10 was found to be 74mM for this system.

Figure 5 shows optimisation of creatine kinase concentration. Optimum concentration was found to be 0.08mg/ml for this system.

Figure 6 shows how % translation could be improved more than two times by using fresh creatine phosphate.

- 15 Figure 7 shows optimisation of creatine phosphate concentration. Optimum concentration was found to be 23mM for this system.

Figure 8 shows optimisation of spermidine concentration. Optimum concentration was found to be 0.1mM for the c-luc-a RNA system.

- Figure 9 shows optimisation of tRNA concentration. Optimum concentration was found
20 to be 100 μ g/ml for the luc RNA system.

Figure 10 shows optimisation of temperature, amino acid concentration and DTT concentration using c-luc-a RNA. Optimum temperature was found to be 25°C. Optimum amino acids concentration was found to be 60 μ M. Optimum DTT concentration was found to be 0mM.

	<u>Range tested</u>	<u>Optimal</u>	<u>RNA</u>	<u>Observations</u>
Creatine phosphate	0 - 168 mM	17 - 23 mM	Luc	Use fresh (thaw only once)
Creatine kinase	0 - 1.68 mg/ml	0.08 mg/ml	Luc	
KOAc	0 - 200 mM	74 mM	Luc	Should be optimised for each RNA template
Mg(OAc) ₂	0 - 3 mM	0.3 mM		
Spermidine	0 - 1.8 mM	0.1 mM	c-Luc-a	0.3 mM or no spermidine also works for some mRNAs
amino acids	0 - 100 μ M	60 μ M	c-Luc-a	
DTT	0 - 4 mM	0 mM	c-Luc-a	Should be optimised for each RNA template
tRNA	0 - 150 μ g/ml	100 μ g/ml	Luc	
Temperature	18 - 37°C	25 °C	c-Luc-a	

Example 3: Comparison between method of the invention and that described in Scott *et al.*, (1979) *Biochemistry*, 18(8): 1588-1594.

Drosophila embryo extracts were prepared in parallel either according to the method described in Scott *et al.* (1979) or to the method of the invention. These extracts were assayed for translation using either the conditions described in Scott *et al.* (1979) or the conditions described above.

Briefly, the Scott protocol consisted in dechorionating the embryos in a solution of 50% ethanol/50% chlorox for 1 m in under agitation, washing the embryos 5 times in PBS after dechorionation using a table-top centrifuge at top speed to obtain the embryo pellet and homogenising the embryo pellet in 10mM HEPES pH7.4, 6mM β -mercaptoethanol.

The extracts obtained using this method were used in an *in vitro* translation reaction using either the conditions described in Example 2.1, those described by Scott *et al* (1979) or those described by Scott with an increased time of translation. The RNA used

for translation encoded the firefly luciferase and contained either a cap (c), a poly(A) (a), or both (c-a). The results (Figures 1 and 2) indicate that by using our extract in our *in vitro* translation conditions a high degree of translation and synergism was obtained. When the Scott extract was used under Scott's *in vitro* translation conditions, no synergism was obtained and a very low level of translation was observed. Using combination of both extracts with either ours or Scott's *in vitro* translation conditions, low levels of translation with various degrees of synergy were obtained. Any apparent synergy in this figure exhibited by the Scott extract using our conditions is believed only to be because the very low level of translation seen gives unreliably low levels of luciferase activity. Accordingly, the (c), (a) and (c-a) figures cannot be compared. These data indicate that the combination of our extract (Example 1) with our *in vitro* translation conditions (Example 2.1) make possible a high degree of translation and synergism.

Comparative results of translation levels and levels of synergy found between the 5' cap and the 3' poly(A) tail of the mRNA template are shown in Figures 1 and 2.

Translation levels using the Scott method were very low. Translation levels using the extract and method of the invention were high and the level of synergy noted was also high (12x and 15x). Figure 2 shows the results of Figure 1 on an expanded scale.

Example 4: Preparation of cell-free extracts from HeLa cells

HeLa cells were maintained at exponential growth in suspension cultures at 37° C in Jocklik's Medium supplemented with 5% newborn bovine serum. Approximately 204 litres of cells at densities of $3-6 \times 10^5$ cells/ml were harvested by centrifugation at 700 g for 15 minutes and washed three times with phosphate buffered saline (PBS) at 4° C.

Pelleted cells were resuspended in 1 volume of ice-cold hypotonic buffer containing Hepes 10 mM, pH 7.6; KOAc 10 mM; Mg(Oac)₂ 0.5 mM; and Dithiothreitol 5 mM. Reduced amounts of DTT are not recommended and addition of protease inhibitors is optional.

After 2-10 minutes on ice, cells were broken with 10-30 strokes of a tight-fitting Dounce homogeniser (pestle type B).

Cell lysates were centrifuged at 14,000 g for 5 minutes (a longer centrifugation is not recommended) at 4° C. Supernatant fluid was divided into aliquots frozen in liquid nitrogen and stored at -80° C. Addition of glycerol is not necessary.

Example 5: In vitro translation assay

- 5 Incubation mixtures contain:
- untreated extract 40% of the final volume
 - Hepes buffer, pH 7.6 16 mM (+4 mM of the lysate)
 - KOAc 50 mM (+4 mM of the lysate)
 - Mg(Oac)₂ 2.5 mM (+0.2 mM of the lysate)
- 10
- amino acid mixture 100 µM
 - spermidine 0.1 mM
 - ATP 0.8 mM
 - GTP 0.1 mM
 - Creatine phosphate 20 µM
- 15
- Creatine kinase 50 ng/µl
 - Calf liver tRNA 100 ng/µl
 - RNA template 1 ng/µl

Reaction mixtures, typically at a final volume of 12.5µl were incubated at 37° C for 90 minutes.

- 20 In order to obtain these optimal values for the translation assay, a range of concentrations of the main components were tested with a capped and polyadenylated mRNA expressing the bacterial enzyme Chloramphenicol Acetyltransferase (CAT). Translational efficiency is judged by the amount of the CAT protein (ng) measured by a colorimetric enzyme immunoassay (CAT ELISA, Böhringer Mannheim).
- 25 Two sets of transcripts coding for the reporter enzymes CAT (Preiss. T. & Hentze. M. W. 1998 Nature 392:516-520) and firefly luciferase. Luc, (Iizuka. N. et al. 1994 Mol. Cell. Biol. 14:7322-7330) were tested in the translation assay and the functional half life of each transcript has been determined to be as follows:
- 5' capped (Cap, m7Gppp) 40 minutes

- 5' capped and polyadenylated (pA, 98 adenines) 55 minutes
- uncapped <10 minutes
- uncapped and polyadenylated <10 minutes

Figure 8 shows a northern blot analysis of the transcripts during the translation assay.

- 5 Bars represent radioactive intensity of each messenger measured by a phosphoimager. Graph (A) shows the four messengers at different times, from 0 to 150 min. Graph (B) shows the Cap and the Cap-pA transcripts from 0 to 90 min.

- The efficiency of translation of capped mRNAs having a polyA tail is thus remarkably higher than the one lacking it. Using several batches of CAT transcripts, it has been
- 10 observed that differences between capped and capped-polyadenylated messengers are in a range of 15 to 60 fold. Uncapped mRNA are poorly translated and are also rapidly degraded.

- In order to study the PolyA effect independently from the presence of the Cap, the stability of the Luc transcripts was enhanced using the cap structure analog
- 15 G(5')ppp(5')A (Acap).

Figure 9 shows the time course of translation of the Luc messengers. Efficiency of translation is represented by light emission measured by a luminometer. Graph (B) shows the data represented in graph (A) with the y axis maximum at 20000 luminescence instead of 550000.

- 20 The functional half life of each transcript has been determined to be :

- Cap: 40 minutes
 - Cap and pA: 40 minutes
 - Acap: 25 minutes
 - Acap and pA: 60 minutes
- 25
- uncapped: 12 minutes
 - uncapped and pA: 20 minutes

The synergism of the polyA and the Cap structures is shown by the 40 fold difference of translation of the Cap-pA mRNA with respect to the sum of the translation of the two mRNAs Cap and Acap-pA.

References

Becker PB, Tsukiyama T and Wu C (1994). Chromatin assembly extracts from *Drosophila* embryos. *Methods Cell Biol.* 44:207-23.

Scott, M. P., Storti, R. V., Pardue, M. L. and Rich A. (1979) Cell-free protein synthesis
5 in lysates of *Drosophila melanogaster* cells. *Biochem.* 18: 1588-1594.

CLAIMS

1. A method for the *in vitro* translation of a ribonucleic acid template, said ribonucleic acid having both a 5' cap and a 3' poly A tail, said method comprising incubating a cell
5 extract of a multicellular eukaryote with said ribonucleic acid template under conditions such that translation of the RNA template to produce its encoded protein by one or more components in the cell extract occurs and the amount of the encoded protein thus produced is greater than the total of (a) the amount of the encoded protein that is produced under said conditions when the ribonucleic acid template has a 5' cap but no
.0 3' poly A tail, plus (b) the amount of the encoded protein that is produced under said conditions when the ribonucleic acid template has a 3' poly A tail but no 5' cap.
2. The method according to claim 1 wherein the cell extract is an animal cell extract.
- 5 3. The method according to claim 2 wherein the cell extract is a mammalian cell extract.
4. The method according to claim 3 wherein the cell extract is a human cell extract.
5. The method according to claim 2 wherein the cell extract is an insect cell extract.
- .0 6. The method according to claim 5 wherein the cell extract is a *Drosophila* cell extract.
7. The method according to claim 6 wherein the cell extract is a *Drosophila* embryo cell
extract.
- 5 8. The method according to claim 7 wherein the *Drosophila* embryo cell extract is prepared by a method comprising dechorionating *Drosophila* embryos in an aqueous isotonic buffer comprising detergent and bleach.
- 0 9. The method according to claim 8 wherein the detergent is Triton X-100 and the bleach is sodium hypochlorite.

10. The method according to either of claims 8 or 9 which further comprises after dechorionating, the steps of washing the embryos; homogenizing the embryos to produce a homogenate; centrifuging the homogenate; and recovering non-pelleted material from the centrifuged homogenate.
- 5
11. The method according to claim 6 wherein the cell extract is a *Drosophila* ovary cell extract.
12. The method according to claim 11 wherein the *Drosophila* ovary cell extract is prepared
- 0 by a method comprising (a) obtaining *Drosophila* ovaries in isotonic medium; (b) washing the ovaries so as to transfer the ovaries slowly from isotonic medium to hypotonic medium, such that the volume of the ovaries increases; (c) homogenising the ovaries to produce a homogenate; (d) centrifuging the homogenate to form an ovary cell extract; and (e) recovering non-pelleted material from the centrifuged homogenate.
- 5
13. The method according to any one of the preceding claims wherein said conditions comprise the presence of the following: creatine phosphate, creatine kinase, potassium and magnesium salts, spermidine, amino acids, a reducing agent, and tRNA.
- 0 14. The method according to any one of the preceding claims wherein said conditions comprise a temperature in the range of 18° to 37°C, and said incubating is for at least 90 minutes.
15. The method according to any one of the preceding claims wherein said incubating is at
- 5 25°C.
16. A method for the preparation of a *Drosophila* embryo extract comprising:
- 10 a) dechorionating *Drosophila* embryos in an aqueous isotonic buffer comprising detergent and bleach;
- b) washing the embryos;
- c) homogenizing the embryos to produce a homogenate;
- d) centrifuging the homogenate; and
- e) recovering non-pelleted material from the centrifuged homogenate.

17. A method for the preparation of a *Drosophila* ovary cell extract comprising (a) obtaining *Drosophila* ovaries in isotonic medium comprising detergent and bleach; (b) washing the ovaries so as to transfer the ovaries slowly from isotonic medium to
5 hypotonic medium, such that the volume of the ovaries increases; (c) homogenising the ovaries to produce a homogenate; (d) centrifuging the homogenate to form an ovary cell extract; and (e) recovering non-pelleted material from the centrifuged homogenate.
18. A method according to either of claims 16 or 17, wherein said detergent is non-ionic.
19. A method according to any one of claims 16-18, wherein said homogenising is
10 performed in the presence of protease inhibitors.
20. A method according to any one of claims 16-19 wherein said buffer is EW buffer (0.7% NaCl, 0.4% Triton X-100).
21. A method according to any one of claims 16-20, wherein said embryos or ovaries are incubated in said isotonic buffer for between 1 and 9 minutes.
- 15 22. A method according to claim 21, wherein said embryos or ovaries are incubated for about 3 minutes.
23. A method according to any one of claims 16-22, wherein said embryos or ovaries are incubated at between 18 and 37°C.
24. A method according to claim 23, wherein said embryos or ovaries are incubated at
20 about 25°C.
25. A method according to any one of claims 16-24 wherein said embryos or ovaries are homogenised at about 4°C.
26. A method according to any one of claims 16-25 additionally comprising the step of adding glycerol to the recovered non-pelleted material to a concentration of about 10%
25 by volume.
27. A method for the preparation of a mammalian cell extract comprising the steps of a) collecting cells by centrifugation; b) washing the cells; c) resuspending the cells; d)

homogenising the cells to produce a homogenate; e) centrifuging the homogenate and f) recovering non-pelleted material from the centrifuged homogenate.

28. A method for the *in vitro* translation of a ribonucleic acid template, said method comprising the steps of:

5 a) adding a ribonucleic acid template to a translation mix in the presence of an embryo extract produced by the method of any one of claims 16-26, an ovary cell extract produced by a method according to any one of claims 17-26, or a mammalian cell extract produced by the method of claim 27 to form a reaction mix,

b) incubating the reaction mix for at least 90 minutes.

10 29. A method according to claim 28, wherein said reaction mix comprises spermidine, amino acids, creatine phosphate, creatine kinase, dithiothreitol (DTT), buffer, $\text{Mg}(\text{OAc})_2$, KOAc, tRNA, embryo or cell extract, and RNA template.

30. A method according to claim 29, wherein said buffer is hypotonic buffer (10mM HEPES pH7.4, 5mM DTT).

15 31. A method according to either of claims 29 or 30, wherein said tRNA is calf liver tRNA.

32. A method according to any of claims 29-31, wherein the concentrations of the components of said reaction mix are as follows: spermidine, about 0.1mM; amino acids, about 60 μM ; DTT, 0-4mM; creatine phosphate, 17-23mM; creatine kinase, 0.08mg/ml; buffer, about 24mM; $\text{Mg}(\text{OAc})_2$, 0-3mM; KOAc, 0-200mM; calf liver tRNA, 0-
0 150 $\mu\text{g}/\text{ml}$; embryo or cell extract, about 40% and RNA template, 1-3ng/ μl .

33. A method according to claim 32, wherein the components of said reaction mix are as follows: spermidine, 0.1mM; amino acids, 60 μM ; DTT, 0-4mM; creatine phosphate, 17-23mM; creatine kinase, 0.08mg/ml; HEPES buffer, 24mM; $\text{Mg}(\text{OAc})_2$, 0-3mM; KOAc, 60mM; calf liver tRNA, 100 $\mu\text{g}/\text{ml}$; embryo or cell extract, 40% and RNA
5 template, 3.2ng/ μl .

34. A method according to any one of claims 28-33, wherein said reaction mix is prepared fresh.

35. A method according to any one of claims 29-34, wherein said creatine phosphate is prepared fresh or is newly thawed.
36. The method according to any one of claims 28-35 wherein said ribonucleic acid
5 template is capped and contains a poly A tail of at least 30 adenine nucleotides in length.
37. A *Drosophila* embryo cell extract, ovary cell extract or mammalian cell extract produced by the method of any one of claims 16-27.
- 10 38. Use of embryo or cell extract according to claim 37 in a method of *in vitro* translation of a ribonucleic acid template.
39. A method according to any one of claims 1-15 wherein said cell extract is prepared according to the method of any one of claims 16 to 27.
40. A method of identifying a candidate molecule that increases or decreases the amount of
15 a produced translation product, comprising:
a) performing the method according to any one of claims 1-15, 28-36 or 39 wherein said conditions or reaction mix comprise the presence of one or more candidate molecules being screened for the ability to increase or decrease the amount of the encoded protein that is produced; and
20 b) identifying any one or more candidate molecules that increase or decrease the amount of the encoded protein that is produced relative to the amount of the encoded protein that is produced in the absence of the one or more candidate molecules.
41. A purified candidate molecule that is identified according to the method of claim 40.
25
42. A kit for *in vitro* translation of ribonucleic acid comprising the following ingredients: spermidine, amino acids, creatine phosphate, creatine kinase, optionally dithiothreitol, buffer, Mg(OAc)₂, KOAc, tRNA, *Drosophila* or mammalian cell extract according to claim 37, and RNA template.
- 30 43. A kit according to claim 42, wherein said buffer is HEPES buffer.


44. A kit according to either of claims 42 or 43, wherein said tRNA is calf liver tRNA.

1/10


FIG. 1**Experiment E-872 Blow up (one set of data eliminated)**


c = cap, a = poly(A), c-a = cap and poly(A)

Synergism = fold over the sum of c+a value




2/10



3/10
FIG. 3

- Luc RNA FROM PROMEGA
- FOR c-luc-a OPTIMAL IS ALSO 0.3mM (THOMAS' DATA)


FIG. 5

- Luc RNA FROM PROMEGA
- MERGE OF 2nd EXP

SUBSTITUTE SHEET (RULE 26)


4/10

FIG. 4



- Luc RNA FROM PROMEGA
- FOR c-luc-a RNA OPTIMAL IS 40mM (THOMAS' DATA)
- MERGE OF 2 ≠ EXP.

FIG. 6




- Luc RNA FROM PROMEGA

SUBSTITUTE SHEET (RULE 26)

5/10

FIG. 7




• Luc RNA FROM PROMEGA

—□— % Translation

• MERGE OF 2 \neq EXP.


FIG. 8



▨ % Translation

SUBSTITUTE SHEET (RULE 26)


6/10



7/10


FIG. 10

E876
c-luc-a RNA



8/10

FIG. 11



9/10

FIG. 12A





FIG. 12B



SUBSTITUTE SHEET (RULE 26)

10/10

FIG. 13A





FIG. 13B



(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
31 August 2000 (31.08.2000)

PCT

(10) International Publication Number
WO 00/50586 A3(51) International Patent Classification⁷: C12N 15/09,
C12P 21/02, C07K 14/435

(21) International Application Number: PCT/IB00/00219

(22) International Filing Date: 22 February 2000 (22.02.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
9904004.0 22 February 1999 (22.02.1999) GB
9919369.0 16 August 1999 (16.08.1999) GB(71) Applicant (for all designated States except US): EURO-
PEAN MOLECULAR BIOLOGY LABORATORY
[DE/DE]; Postfach 10 22 09, Meyerhofstrasse 1, D-69012
Heidelberg (DE).

(72) Inventors; and

(75) Inventors/Applicants (for US only): GEBAUER, Fa-
tima [ES/DE]; Lerchenstrasse 28, D-69251 Gaiberg
(DE). CORONA, Davide [IT/DE]; Brechtelstrasse 8-10,
D-69126 Heidelberg (DE). BERGAMINI, Giovanna
[IT/DE]; Römerstrasse 25, D-69115 Heidelberg (DE).
BECKER, Peter [DE/DE]; Salzstrasse 9, D-82194
Gröbenzell (DE). HENTZE, Matthias [DE/DE]; Meri-
anstrasse 5, D-69117 Heidelberg (DE).(74) Agent: HALLYBONE, Huw, George; Carpmaels &
Ransford, 43 Bloomsbury Square, London WC1A 2RA
(GB).(81) Designated States (national): AE, AL, AM, AT, AU, AZ,
BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK,
DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL,
IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU,
LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT,


[Continued on next page]

(54) Title: IN VITRO TRANSLATION SYSTEM

Experiment E-872 Blow up (one set of data eliminated)

c = cap, a = poly(A), c-a = cap and poly(A)

Synergism = fold over the sum of c+a value

(57) Abstract: The invention relates to a cell-free translation system using cell extracts, that reproduces the synergism that occurs during translation *in vivo* between the *cap* and the *poly(A)* tail structures of an mRNA molecule. Using this system, the amount of encoded protein produced from translation of an mRNA molecule is greater than the total of (a) the amount of the encoded protein that is produced under the same conditions when the ribonucleic acid template has a 5' cap but no 3' poly A tail, plus (b) the amount of the encoded protein that is produced under the same conditions when the ribonucleic acid template has a 3' poly A tail but no 5' cap.

WO 00/50586 A3



RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA,
UG, US, UZ, VN, YU, ZA, ZW.

Published:

— With international search report.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(88) Date of publication of the international search report:
4 January 2001

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/IB 00/00219

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N15/09 C12P21/02 C07K14/435

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12P C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	SCOTT M P ET AL.: "Cell-free protein synthesis in lysates of Drosophila melanogaster cells" BIOCHEMISTRY, vol. 18, no. 8, 1979, pages 1588-1594, XP002140264 cited in the application. the whole document	1,2,5-7, 11,13, 40,42,43
X	--- HUSSAIN I AND LEIBOWITZ M J: "Translation of homologous and heterologous messenger RNAs in a yeast cell-free system" GENE, vol. 46, 1986, pages 13-23, XP002140265 the whole document --- -/-	1,13,14, 28,29, 34-36, 40,42,43

☒ Further documents are listed in the continuation of box C.☐ Patent family members are listed in annex.

* Special categories of cited documents:

A document defining the general state of the art which is not considered to be of particular relevance

E earlier document but published on or after the international filing date

L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

O document referring to an oral disclosure, use, exhibition or other means

P document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

Z document member of the same patent family

Date of the actual completion of the international search

15 June 2000

Date of mailing of the international search report

31. 08. 2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Oderwald, H

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	IIZUKA N ET AL.: "Cap-dependent and cap-independent translation by internal initiation of mRNAs in cell extracts prepared from <i>Saccharomyces cerevisiae</i> " MOLECULAR AND CELLULAR BIOLOGY, vol. 14, no. 11, November 1994 (1994-11), pages 7322-7330, XP000914650 cited in the application the whole document	1,40
P,X	--- GEBAUER F ET AL.: "Translational control of dosage compensation in <i>Drosophila</i> by Sex-lethal: cooperative silencing via the 5' and 3' UTRs of msl-2 mRNA is independent of the poly(A) tail" EMBO JOURNAL, vol. 18, no. 21, November 1999 (1999-11), pages 6146-6154, XP002140266 the whole document -----	1,2, 5-10, 13-15, 28-36, 39,40, 42-44

INTERNATIONAL SEARCH REPORT

International application No.
PCT/IB 00/00219

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. ☒ Claims Nos.: 41
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
see FURTHER INFORMATION sheet PCT/ISA/210

3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.

2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.

3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:

4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:
1-15, 28-36, 39, 40, 42-44

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Claims Nos.: 41

~~Claim 41~~ refers to an inhibitor or acticator of translation without giving a true technical characterization. Moreover, no such compounds are defined in the application. In consequence, the scope of said claims is ambiguous and vague, and their subject-matter is not sufficiently disclosed and supported (Art. 5 and 6 PCT). No search can be carried out for such purely speculative claims whose wording is, in fact, a mere recitation of the results to be achieved.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

1. Claims: 1-15, 28-36, 39, 40, 42-44

Method for in vitro translation of a ribonucleic acid template, a method of identifying a candidate molecule that increases or decreases the amount of a produced translation product, a kit for in vitro translation.

2. Claims: 16-27, 37, 38

Methods for the preparation of a Drosophila embryo extract, a Drosophila ovary cell extract and a mammalian cell extract, said cell extracts and its use in in vitro translation.